

**MTN 015 HIV-1 Seroconverter Study**  
**Data as of 09JUN2020**  
**Visit Adherence: Procedure Completion, by Site**

	<i>All Sites</i>	<i>Blantyre, Malawi</i>	<i>Lilongwe, Malawi</i>	<i>Lusaka, Zambia</i>	<i>Spilhaus/ Zimbabwe</i>	<i>Seke South/ Zimbabwe</i>	<i>MRC - Chatsworth/ RK Khan</i>
Visits at which lab tests are required	3527	259	221	124	186	128	478
T Cell Subsets Completed	3508 (99.5%)	259 (100.0%)	219 (99.1%)	124 (100.0%)	185 (99.5%)	127 (99.2%)	478 (100.0%)
HIV RNA Completed	3498 (99.2%)	259 (100.0%)	220 (99.5%)	124 (100.0%)	186 (100.0%)	127 (99.2%)	478 (100.0%)
Visits at which Plasma and Vaginal Swab collection are required	3527	259	221	124	186	128	478
Plasma Collection Completed	3494 (99.1%)	255 (98.5%)	217 (98.2%)	124 (100.0%)	186 (100.0%)	128 (100.0%)	473 (99.0%)
Vaginal Swab Collection Completed	3214 (91.1%)	255 (98.5%)	208 (94.1%)	120 (96.8%)	181 (97.3%)	124 (96.9%)	439 (91.8%)
Visits at which PBMC collection are required [1]	3114	244	207	-	176	122	462
PBMC Collection Completed	2884 (92.6%)	131 (53.7%)	131 (63.3%)	-	175 (99.4%)	120 (98.4%)	460 (99.6%)
Visits at which Cervicovaginal Lavage collection are required	3279	229	196	117	169	118	454
Cervicovaginal Lavage Collection Completed	3028 (92.3%)	228 (99.6%)	189 (96.4%)	114 (97.4%)	168 (99.4%)	117 (99.2%)	410 (90.3%)
Visits at which Serum collection are required	487	27	21	14	19	17	74
Serum Collection Completed	362 (74.3%)	22 (81.5%)	14 (66.7%)	14 (100.0%)	11 (57.9%)	11 (64.7%)	62 (83.8%)
Visits at which STD tests are required	1611	117	95	60	75	57	236
Vaginal pH Completed	1453 (90.2%)	117 (100.0%)	77 (81.1%)	58 (96.7%)	71 (94.7%)	53 (93.0%)	226 (95.8%)
Syphilis Tests Completed	1545 (95.9%)	117 (100.0%)	93 (97.9%)	59 (98.3%)	71 (94.7%)	54 (94.7%)	232 (98.3%)
N. gonorrhoea Tests Completed	1549 (96.2%)	117 (100.0%)	93 (97.9%)	59 (98.3%)	72 (96.0%)	54 (94.7%)	234 (99.2%)
C. trachomatis Tests Completed	1548 (96.1%)	117 (100.0%)	93 (97.9%)	59 (98.3%)	72 (96.0%)	54 (94.7%)	234 (99.2%)

[1] PBMC collection is required only at sites with capacity. These sites do not have capacity: Lusaka, Zambia; and Kampala, Uganda.

**MTN 015 HIV-1 Seroconverter Study**  
**Data as of 09JUN2020**  
**Visit Adherence: Procedure Completion, by Site**

	<i>All Sites</i>	<i>MRC - Botha's Hill</i>	<i>MRC - Umkomaas</i>	<i>Kampala, Uganda</i>	<i>Zengeza/ Zimbabwe</i>	<i>MRC - Overport</i>	<i>MRC - Isipingo</i>
Visits at which lab tests are required	3527	178	188	158	106	118	202
T Cell Subsets Completed	3508 (99.5%)	177 (99.4%)	188 (100.0%)	158 (100.0%)	106 (100.0%)	118 (100.0%)	201 (99.5%)
HIV RNA Completed	3498 (99.2%)	174 (97.8%)	187 (99.5%)	158 (100.0%)	105 (99.1%)	118 (100.0%)	201 (99.5%)
Visits at which Plasma and Vaginal Swab collection are required	3527	178	188	158	106	118	202
Plasma Collection Completed	3494 (99.1%)	178 (100.0%)	188 (100.0%)	158 (100.0%)	106 (100.0%)	116 (98.3%)	201 (99.5%)
Vaginal Swab Collection Completed	3214 (91.1%)	172 (96.6%)	173 (92.0%)	139 (88.0%)	99 (93.4%)	96 (81.4%)	189 (93.6%)
Visits at which PBMC collection are required [1]	3114	177	183	-	96	115	198
PBMC Collection Completed	2884 (92.6%)	177 (100.0%)	182 (99.5%)	-	96 (100.0%)	113 (98.3%)	197 (99.5%)
Visits at which Cervicovaginal Lavage collection are required	3279	177	181	141	89	112	196
Cervicovaginal Lavage Collection Completed	3028 (92.3%)	171 (96.6%)	165 (91.2%)	133 (94.3%)	88 (98.9%)	92 (82.1%)	186 (94.9%)
Visits at which Serum collection are required	487	35	31	19	13	20	37
Serum Collection Completed	362 (74.3%)	28 (80.0%)	26 (83.9%)	11 (57.9%)	6 (46.2%)	19 (95.0%)	32 (86.5%)
Visits at which STD tests are required	1611	91	85	67	41	53	93
Vaginal pH Completed	1453 (90.2%)	88 (96.7%)	74 (87.1%)	63 (94.0%)	35 (85.4%)	42 (79.2%)	81 (87.1%)
Syphilis Tests Completed	1545 (95.9%)	89 (97.8%)	83 (97.6%)	66 (98.5%)	37 (90.2%)	53 (100.0%)	91 (97.8%)
N. gonorrhoea Tests Completed	1549 (96.2%)	87 (95.6%)	83 (97.6%)	66 (98.5%)	37 (90.2%)	53 (100.0%)	87 (93.5%)
C. trachomatis Tests Completed	1548 (96.1%)	87 (95.6%)	83 (97.6%)	66 (98.5%)	37 (90.2%)	53 (100.0%)	87 (93.5%)

[1] PBMC collection is required only at sites with capacity. These sites do not have capacity: Lusaka, Zambia; and Kampala, Uganda.

**MTN 015 HIV-1 Seroconverter Study**  
**Data as of 09JUN2020**  
**Visit Adherence: Procedure Completion, by Site**

	<i>All Sites</i>	<i>MRC - Tongaat</i>	<i>MRC - Verulam</i>	<i>Aurum</i>	<i>eThekwini</i>	<i>WRHI</i>	<i>Soweto MTN/ PHRU</i>
Visits at which lab tests are required	3527	333	211	52	394	157	34
T Cell Subsets Completed	3508 (99.5%)	331 (99.4%)	201 (95.3%)	52 (100.0%)	394 (100.0%)	156 (99.4%)	34 (100.0%)
HIV RNA Completed	3498 (99.2%)	328 (98.5%)	197 (93.4%)	52 (100.0%)	394 (100.0%)	156 (99.4%)	34 (100.0%)
Visits at which Plasma and Vaginal Swab collection are required	3527	333	211	52	394	157	34
Plasma Collection Completed	3494 (99.1%)	331 (99.4%)	201 (95.3%)	52 (100.0%)	390 (99.0%)	156 (99.4%)	34 (100.0%)
Vaginal Swab Collection Completed	3214 (91.1%)	278 (83.5%)	160 (75.8%)	49 (94.2%)	357 (90.6%)	144 (91.7%)	31 (91.2%)
Visits at which PBMC collection are required [1]	3114	326	204	50	370	151	33
PBMC Collection Completed	2884 (92.6%)	324 (99.4%)	194 (95.1%)	50 (100.0%)	370 (100.0%)	131 (86.8%)	33 (100.0%)
Visits at which Cervicovaginal Lavage collection are required	3279	323	199	50	347	148	33
Cervicovaginal Lavage Collection Completed	3028 (92.3%)	273 (84.5%)	151 (75.9%)	47 (94.0%)	329 (94.8%)	139 (93.9%)	28 (84.8%)
Visits at which Serum collection are required	487	51	28	12	41	20	8
Serum Collection Completed	362 (74.3%)	41 (80.4%)	18 (64.3%)	12 (100.0%)	15 (36.6%)	12 (60.0%)	8 (100.0%)
Visits at which STD tests are required	1611	165	100	28	160	69	19
Vaginal pH Completed	1453 (90.2%)	138 (83.6%)	77 (77.0%)	26 (92.9%)	148 (92.5%)	62 (89.9%)	17 (89.5%)
Syphilis Tests Completed	1545 (95.9%)	147 (89.1%)	92 (92.0%)	27 (96.4%)	152 (95.0%)	65 (94.2%)	17 (89.5%)
N. gonorrhoea Tests Completed	1549 (96.2%)	143 (86.7%)	94 (94.0%)	27 (96.4%)	157 (98.1%)	67 (97.1%)	19 (100.0%)
C. trachomatis Tests Completed	1548 (96.1%)	142 (86.1%)	94 (94.0%)	27 (96.4%)	157 (98.1%)	67 (97.1%)	19 (100.0%)

[1] PBMC collection is required only at sites with capacity. These sites do not have capacity: Lusaka, Zambia; and Kampala, Uganda.