

**MTN 020 - A Study to Prevent Infection with a Ring for Extended Use (ASPIRE)**  
**Data as of September 21, 2015**

**Baseline Contraceptive Use by Site**

	All Sites	Malawi - Blantyre	Malawi - Lilongwe	SA - Cape Town	SA - CAPRISA eThekweni	SA - MRC/ Botha's Hill	SA - MRC/ Chatsworth	SA - MRC/ Isipingo	SA - MRC/ Tongaat
Participants Enrolled	2629	130	142	166	244	180	150	117	103
Participants with Baseline Family Planning Form	2629	130	142	166	244	180	150	117	103
<b>Current Method of Contraception<sup>1</sup></b>									
None	0	0	0	0	0	0	0	0	0
Spermicide	0	0	0	0	0	0	0	0	0
Diaphragm	0	0	0	0	0	0	0	0	0
Sponge	0	0	0	0	0	0	0	0	0
Intrauterine device	325 (12.4%)	0	6 (4.2%)	18 (10.8%)	32 (13.1%)	1 (0.6%)	2 (1.3%)	1 (0.9%)	0
Oral contraceptives	287 (10.9%)	6 (4.6%)	0	6 (3.6%)	19 (7.8%)	29 (16.1%)	29 (19.3%)	12 (10.3%)	30 (29.1%)
Injectable contraceptives	1452 (55.2%)	64 (49.2%)	51 (35.9%)	138 (83.1%)	176 (72.1%)	148 (82.2%)	112 (74.7%)	99 (84.6%)	65 (63.1%)
Depo	1070 (40.7%)	64 (49.2%)	51 (35.9%)	102 (61.4%)	114 (46.7%)	114 (63.3%)	78 (52.0%)	72 (61.5%)	37 (35.9%)
Depo subq	1 (<0.1%)	0	0	0	0	0	0	1 (0.9%)	0
NET-EN	381 (14.5%)	0	0	36 (21.7%)	62 (25.4%)	34 (18.9%)	34 (22.7%)	26 (22.2%)	28 (27.2%)
Cyclofem	0	0	0	0	0	0	0	0	0
Other	0	0	0	0	0	0	0	0	0
Patch	0	0	0	0	0	0	0	0	0
Implants	501 (19.1%)	41 (31.5%)	79 (55.6%)	2 (1.2%)	12 (4.9%)	0	2 (1.3%)	0	0
Female condoms	1 (<0.1%)	0	0	0	1 (0.4%)	0	0	0	0
Natural methods	0	0	0	0	0	0	0	0	0
Male condoms	96 (3.7%)	0	0	48 (28.9%)	20 (8.2%)	0	1 (0.7%)	0	0
Sterilization	78 (3.0%)	19 (14.6%)	6 (4.2%)	5 (3.0%)	6 (2.5%)	2 (1.1%)	5 (3.3%)	5 (4.3%)	8 (7.8%)
Partner has a vasectomy	0	0	0	0	0	0	0	0	0
Other	0	0	0	0	0	0	0	0	0

<sup>1</sup> Participants could report more than one method of contraception.

**MTN 020 - A Study to Prevent Infection with a Ring for Extended Use (ASPIRE)**  
**Data as of September 21, 2015**

**Baseline Contraceptive Use by Site Continued**

	All Sites	SA - MRC/ Verulam	SA - MRC/ Umkomaas	SA - WRHI	Uganda - Kampala	Zimbabwe - Seke South	Zimbabwe - Spilhaus	Zimbabwe - Zengeza
Participants Enrolled	2629	150	103	213	253	224	230	224
Participants with Baseline Family Planning Form	2629	150	103	213	253	224	230	224
<b>Current Method of Contraception<sup>1</sup></b>								
None	0	0	0	0	0	0	0	0
Spermicide	0	0	0	0	0	0	0	0
Diaphragm	0	0	0	0	0	0	0	0
Sponge	0	0	0	0	0	0	0	0
Intrauterine device	325 (12.4%)	3 (2.0%)	0	22 (10.3%)	71 (28.1%)	77 (34.4%)	54 (23.5%)	38 (17.0%)
Oral contraceptives	287 (10.9%)	38 (25.3%)	18 (17.5%)	60 (28.2%)	14 (5.5%)	10 (4.5%)	8 (3.5%)	8 (3.6%)
Injectable contraceptives	1452 (55.2%)	100 (66.7%)	82 (79.6%)	126 (59.2%)	112 (44.3%)	52 (23.2%)	59 (25.7%)	68 (30.4%)
Depo	1070 (40.7%)	59 (39.3%)	60 (58.3%)	28 (13.1%)	112 (44.3%)	52 (23.2%)	59 (25.7%)	68 (30.4%)
Depo subq	1 (<0.1%)	0	0	0	0	0	0	0
NET-EN	381 (14.5%)	41 (27.3%)	22 (21.4%)	98 (46.0%)	0	0	0	0
Cyclofem	0	0	0	0	0	0	0	0
Other	0	0	0	0	0	0	0	0
Patch	0	0	0	0	0	0	0	0
Implants	501 (19.1%)	0	0	5 (2.3%)	49 (19.4%)	85 (37.9%)	110 (47.8%)	116 (51.8%)
Female condoms	1 (<0.1%)	0	0	0	0	0	0	0
Natural methods	0	0	0	0	0	0	0	0
Male condoms	96 (3.7%)	9 (6.0%)	0	14 (6.6%)	0	0	0	4 (1.8%)
Sterilization	78 (3.0%)	9 (6.0%)	3 (2.9%)	0	7 (2.8%)	0	1 (0.4%)	2 (0.9%)
Partner has a vasectomy	0	0	0	0	0	0	0	0
Other	0	0	0	0	0	0	0	0

<sup>1</sup> Participants could report more than one method of contraception.