

MTN 034 - Reversing the Epidemic in Africa with Choices in HIV Prevention (REACH)
Data as of December 5, 2022

Baseline Contraceptive Use by Site

	All Sites	SA - Cape Town	SA - Johannesburg	Uganda - Kampala	Zimbabwe - Spilhaus
Participants Enrolled	247	60	67	60	60
Participants with Family Planning Log	247	60	67	60	60
Method of Contraception¹					
Spermicide	0	0	0	0	0
Sponge	0	0	0	0	0
Oral contraceptive birth control pills (Ortho Evra) - The Patch	0	0	0	0	0
Implants	111 (44.9%)	11 (18.3%)	30 (44.8%)	37 (61.7%)	33 (55.0%)
Female condoms	0	0	0	0	0
Male condoms	0	0	0	0	0
Sterilization	0	0	0	0	0
Diaphragm	0	0	0	0	0
Copper IUD	26 (10.5%)	1 (1.7%)	7 (10.4%)	9 (15.0%)	9 (15.0%)
Hormonal IUD (e.g., Mirena, Lilette, Skyla)	1 (0.4%)	0	1 (1.5%)	0	0
Injectable contraceptive - Depo	65 (26.3%)	20 (33.3%)	15 (22.4%)	12 (20.0%)	18 (30.0%)
Injectable contraceptive - NET-EN	42 (17.0%)	28 (46.7%)	14 (20.9%)	0	0
Injectable contraceptive - Cyclofem	0	0	0	0	0
Injectable contraceptive - Other	2 (0.8%)	0	0	2 (3.3%)	0
Natural methods such as the withdrawal or rhythm method	0	0	0	0	0
Sex with partner who had vasectomy	0	0	0	0	0
Emergency contraception	0	0	0	0	0
Other	0	0	0	0	0
Missing Family Planning CRF	0	0	0	0	0

¹ Participants could report more than one method of contraception.